

13th Intern. Conf. on the Applic. of Density Functional Theory in
Chemistry and Physics, DFT09, Lyon, France.

QM/MM Meta-Dynamics Study of Asparagine Deamidation in Proteins

Saron Catak^{a,*}, Bart De Sterck^a, Ewald Pauwels^a, Gerald Monard^b, Manuel F. Ruiz-López^b, Viktorya Aviyente^c, Michel Waroquier^a, Veronique Van Speybroeck^{a,*}

^aCenter for Molecular Modeling, Ghent University, Proeftuinstraat 86, B-9000 Gent, Belgium


^bEquipe de Chimie et Biochimie Théoriques, UMR CNRS-UHP No. 7565, Nancy-Université, BP 239, 54506 Vandoeuvre-lès-Nancy, France

^cDepartment of Chemistry, Bogazici University, 34342 Bebek, Istanbul, Turkey

* Corresponding authors: saron.catak@ugent.be, veronique.vanspeybroeck@ugent.be

Asparagine (Asn) residues in peptides and proteins are subject to spontaneous non-enzymatic deamidation to form aspartate or iso-aspartate at a rate determined by protein structure and environment [1]. Deamidation causes time-dependent changes in charge and conformation and limits the lifetime of peptides and proteins. This has led to the ‘*molecular clocks*’ hypothesis [2], which suggests deamidation is a biological molecular timing mechanism that could be set to any desired time interval by genetic control of the primary, secondary, and tertiary structure surrounding the amide. A notable example is Bcl-x_L, an anti-apoptotic protein that regulates cell death and undergoes a ‘complete loss’ of function upon deamidation; suppression of deamidation in Bcl-x_L plays a pivotal role in the survival of tumor cells [1].

Deamidation occurs over a wide range of biologically relevant time intervals; this suggests that different mechanisms may be operative. While a succinimide-mediated deamidation pathway was originally suggested [3-5], two new ‘competing’ routes for deamidation of asparagine residues will be presented. In order to compare relative energetics and feasibilities of plausible routes, QM/MM molecular dynamics simulations, in a periodic water box, were performed on a model peptide. Free energies of competing pathways were calculated *via* meta-dynamics calculations utilizing the CP2K code [6], where the catalytic effect of water molecules was explicitly analyzed. These calculations have served as a stepping stone for QM/MM metadynamics calculations on deamidation of Bcl-x_L.



Keywords: Deamidation, asparagine, Bcl-x_L, meta-dynamics, QM/MM, CP2K

[1] Weintraub, S.J. and B.E. Deverman, *Sci. STKE.*, **409**, re7 (2007)

[2] N. E. Robinson and A. B. Robinson, *Proc. Natl. Acad. Sci. USA*, **98**, 944 (2001)

[3] Kim, E., Lowenson, J. D., MacLaren, D. C., Clarke, S., *Proc. Natl. Acad. Sci. USA*, **94**, 6132 (1997)

[4] S. Catak, G. Monard, V. Aviyente and M. F. Ruiz-López, *J. Phys. Chem. A*, **110**, 8354 (2006)

[5] S. Catak, G. Monard, V. Aviyente and M. F. Ruiz-López, *J. Phys. Chem. A*, **113**, 1111 (2009)

[6] <http://cp2k.berlios.de>